

Airway Patency Assessment



Root Cause Resolutions

Dental + Medical Integration

Liz Laney, BSDH, RDH

Oral-Systemic RDH & Educator

This Airway Patency Assessment was created to summarize three common clinical screening tools for airway obstruction risks, which provide a standard protocol for healthcare and dental providers and improves whole-body patient care.

The information within this packet is validated and proven by years of research and testing for identification of approximate airway obstruction at the hypopharyngeal level and has been simplified down to easy-to-use, single-page summaries.

- Modified Mallampati Score
- Friedman Tongue Position
- Friedman Tonsil Grading
- Airway Patency Assessment - summary
- Airway Screening Form - update yearly for each patient
- Intervention Protocols

Please note, the information in this packet is not intended to take the place of talking with a medical provider and should not be used for diagnosis. This was created as a screening guide only. Regardless of the results, if a patient or provider has concerns about noted symptoms, they are encouraged to discuss them further with a medical and/or dental provider.

This packet includes affiliate and partner links for products that I love and recommend daily to my patients. If you decide to make a purchase via the link(s), I may earn a commission at no additional cost to you.

Copyright notice: The information curated in this document is free for personal use only and is protected by copyright laws. No changes may be made, and resale of digital download item(s) is not allowed for personal or commercial use.

Modified Mallampati Score


The original Mallampati Score was created in 1985 as a way to assess and describe the relative size of the tongue compared to the oropharyngeal opening to predict difficult intubation of patients undergoing general anesthesia or entering an operating room or critical care setting.⁵

- The original Mallampati Score was based on a range from Class I-III

The Modified Mallampati Score emerged in 1987 with a range from Class I-IV and was updated again in 1998 to include Class 0-IV, which provides the most detailed breakdown of airway patency and is the version most used in modern medicine.

Guidelines:

- Patient is seated upright
- Assessed when the tongue is maximally protruded
- Classically, this exam is completed without phonation (no vocal sounds made)



In terms of airway health, a Mallampati Class III or Class IV indicates an increased risk of airway blockage during sleep, resulting in obstructive sleep apnea. Because of this, airway providers often use the Modified Mallampati grading scale as a simple guideline for identifying patients who may need further airway interventions, such as myofunctional therapy, a CPAP/BiPAP machine, or possibly even surgical interventions.

As the Modified Mallampati grading scale is not intended to diagnose sleep apnea, it is designed to be used as a screening tool, and patients should seek further assistance about any questions or concerns they have to receive treatment based on their individual needs.




Friedman Tongue Position

The Friedman Tongue Position (FTP) was initially presented in 1999 as a way to evaluate the tongue's position relative to the tonsils or pillars, hard palate, uvula, and soft palate and to approximate airway obstruction at the hypopharyngeal level.³ Comparative to the Modified Mallampati, the FTP is assessed while the tongue is in a neutral position.

Guidelines:

- Patient is seated upright
- Assessed when the tongue is in its natural resting position inside the mouth (not protruded)
 - Have patient open wide five times to establish an average neutral tongue position
 - While breathing normally, assess the base of the tongue relative to the tonsils/tonsillar pillars, hard palate, uvula, and soft palate
- Includes a scale of Grade I-IV
 - Sub-grading of IIa and IIb provide greater descriptions of airway patency



While both the Modified Mallampati score and the Friedman Tongue Position describe the relative position of the base of the tongue to the oropharyngeal opening, the Friedman Tongue Position provides a more detailed evaluation of the tongue's neutral resting position, which is generally more helpful for assessing and identifying potential airway restrictions.

To keep interdisciplinary care teams from misunderstanding or misinterpreting reports from other providers, it's essential that each provider identify which grading scale they utilize, either Modified Mallampati or FTP.



Friedman Tonsil Grading

The Friedman Tonsil Grading system was created to identify enlargement, or hypertrophy, of the palatine tonsils, which are located in the lateral walls of the oropharynx between the palatoglossal arch (anterior) and palatopharyngeal arch (posterior).⁴ The palatal tonsils are part of the lymphatic system and important for both innate and acquired immunity.

Cycle of Dysfunction

- When palatine tonsillar hypertrophy is present, individuals often resort to mouth breathing, which allows harmful pathogens, bacteria, and viruses to enter the body and bypass filtration within the nasal cavity.
- This creates a perpetual immune response and chronic tonsillar hypertrophy, and an individual is unable to restore nasal breathing habits effectively.
- Chronic mouth breathing habits increases an individuals risk of oral microbiome dysbiosis and in turn increases their risk for gingival inflammation (i.e. gingivitis and periodontal disease) and dental caries (i.e. cavities).

A quick and easy chairside assessment to evaluate for possible airway resistance should include palatine tonsil grading.

- Following assessment, should tonsillar hypertrophy be present, intervention protocols should be followed. This gives the greatest chance of lowering inflammation of the tonsillar tissue and restoring airway patency and, therefore, nasal breathing.




Grade 0: absence of tonsillar tissue
Grade 1: tonsils contained within the pillars
Grade 2: tonsils extend to the pillars
Grade 3: tonsils extend beyond the pillars but do not meet at the midline
Grade 4: tonsils extend to the midline




Airway Patency Assessment


Modified Mallampati



Friedman Tongue Position



Friedman Tonsil Grading



Airway Screening Assessment

Patient Name: _____ Date: _____

Patient-Reported Symptoms

- | | |
|---|--|
| <input type="checkbox"/> Snoring | <input type="checkbox"/> Sleep talking, sleepwalking |
| <input type="checkbox"/> Fatigue | <input type="checkbox"/> Night terrors |
| <input type="checkbox"/> Excessive daytime sleepiness | <input type="checkbox"/> Nursing difficulties |
| <input type="checkbox"/> Headaches | <input type="checkbox"/> Irritability |
| <input type="checkbox"/> Neck tension | <input type="checkbox"/> Sleep paralysis |
| <input type="checkbox"/> Difficulty concentrating | <input type="checkbox"/> Drooling |
| <input type="checkbox"/> ADHD | <input type="checkbox"/> Anxiety & depression |
| <input type="checkbox"/> Bedwetting | <input type="checkbox"/> Thumb/digit sucking |
| <input type="checkbox"/> Acid reflux | <input type="checkbox"/> Mouth breathing |

Clinical Presentation

- | | |
|--|---|
| <input type="checkbox"/> Tooth decay | <input type="checkbox"/> Clenching and grinding |
| <input type="checkbox"/> Narrow palate | <input type="checkbox"/> Recession |
| <input type="checkbox"/> Orthodontic relapse | <input type="checkbox"/> Abfraction |
| <input type="checkbox"/> Forward head posture | <input type="checkbox"/> Occlusal wear |
| <input type="checkbox"/> Class II or Class III occlusion | <input type="checkbox"/> Craze lines |
| <input type="checkbox"/> Tongue thrust | <input type="checkbox"/> Cracked teeth |
| <input type="checkbox"/> Anterior or posterior open bite | <input type="checkbox"/> Enlarged tonsils |
| <input type="checkbox"/> Palatal an/or mandibular tori | <input type="checkbox"/> Persistent gingival inflammation |
| <input type="checkbox"/> Teeth crowding | <input type="checkbox"/> TMJ dysfunction |
| <input type="checkbox"/> Erosion - acid reflux | <input type="checkbox"/> Click, crepitus |
| <input type="checkbox"/> Limited tongue space | <input type="checkbox"/> Pain |
| <input type="checkbox"/> Scalloped lateral borders | <input type="checkbox"/> Lock jaw |
| <input type="checkbox"/> Cracked, dry lips | <input type="checkbox"/> Mouth breathing |

Airway Anatomy Evaluation

- | | |
|---|--|
| <input type="checkbox"/> Palatine tonsils - enlarged | <input type="checkbox"/> Tongue function |
| <input type="checkbox"/> Adenoids - enlarged (via CBCT) | <input type="checkbox"/> Tongue Range of Motion Ratio (TRMR) |
| <input type="checkbox"/> Labial frenulum | <input type="checkbox"/> Maxilla |
| <input type="checkbox"/> Inferior turbinate hypertrophy | <input type="checkbox"/> Intermolar width |
| <input type="checkbox"/> Narrow nasal valve(s) | <input type="checkbox"/> Signs of deficient growth |
| <input type="checkbox"/> Tongue to airway position | <input type="checkbox"/> Crowding |
| <input type="checkbox"/> Modified Mallampati | <input type="checkbox"/> Impacted teeth |
| <input type="checkbox"/> Friedman Tongue Position | <input type="checkbox"/> Delayed tooth eruption |
| | <input type="checkbox"/> Retrusive mandible |



Intervention Protocols

Nasal Clearing

One of the top recommended products for nasal clearing is Xlear® Nasal Spray, a natural saline solution with xylitol that inhibits bacteria adhesion and replication within the nasopharynx and promotes a decrease in tonsillar inflammation.



Order Xlear® Nasal Spray



Oral Microbiome Dysbiosis

Dysbiosis, or an imbalance of the good and bad bacteria, is a side effect of mouth breathing, and each person has individual needs for rebalancing their oral microbiome. Common signs of oral dysbiosis includes halitosis (i.e. bad breath), high cavity rate, dry mouth, and gingival inflammation such as gingivitis or periodontal disease.

Use of the **Bristle Oral Health Test** -- an at-home oral microbiome test -- will provide individualized product recommendations and is a great place to start the process of repairing the oral microbiome.

Please note, use of this product does not replace regular visits with medical and/or dental providers.

As a Bristle Partner, I am happy to offer 10% off their Oral Health Test and Oral Probiotics.

Scan the QR code to order!



Dental providers looking to join the Bristle Partner Program can find out more [here](#).



Sources

1. Friedman M, Hwang MS. Evaluation of the patient with obstructive sleep apnea: Friedman tongue position and staging. *Operative Techniques in Otolaryngology-Head and Neck Surgery*. 2015;26(2):85-89. doi:10.1016/j.otot.2015.03.007
2. Friedman M, Salapatras AM, Bonzelaar LB. Updated Friedman Staging System for Obstructive Sleep Apnea. *Adv Otorhinolaryngol*. 2017;80:41-48. doi:10.1159/000470859
3. Friedman tongue position and the staging of obstructive sleep apnea/hypopnea syndrome. *Pocket Dentistry*. July 24, 2016. <https://pocketdentistry.com/friedman-tongue-position-and-the-staging-of-obstructive-sleep-apneahypopnea-syndrome/>.
4. Meegalla N, Downs BW. Anatomy, Head and Neck, Palatine Tonsil (Faucial Tonsils) [Updated 2023 Jun 5]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2024 Jan-. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK538296/>
5. Stutz EW, Rondeau B. Mallampati Score. [Updated 2023 Aug 5]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2024 Jan-. Available from: Stutz EW, Rondeau B. Mallampati Score. [Updated 2023 Aug 5]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2024 Jan-. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK585119/>
6. Thomas M, Saldanha NM. Comparison of upper lip bite test with modified Mallampati score in predicting difficult intubation. *International Journal of Science and Healthcare Research*. 2022;7(1):1-8. doi:10.52403/ijshr.20220101

